

How to Manage Common Conditions for People with Learning Disabilities

A faint, stylized line drawing in the background shows three people sitting at a table. On the table are various items including a bowl, a glass, and some food. The drawing is composed of simple outlines with some color accents like green for foliage and yellow for food items.

This comprehensive guide provides essential information for caregivers and healthcare staff supporting adults with learning disabilities. We'll explore the management of four critical conditions that frequently affect this vulnerable population: diarrhoea, constipation, reflux, and dysphagia. Understanding these conditions and their proper management is crucial for maintaining the health, dignity, and quality of life for those in your care.


Table of Contents	Pg	Video m:sec
2. Understanding Diarrhoea in Learning Disabilities	3	0:29
3. Causes and Initial Management of Diarrhoea	4	1:03
4. Dietary Management Strategies	5	3:55
5. Understanding Constipation: A Critical Concern	6	7:55
6. Recognising and Responding to Constipation	7	8:50
7. Prevention Strategies for Constipation	8	11:05
8. Managing Reflux in Learning Disabilities	9	14:55
9. Preventing Reflux in Individuals with Learning Disabilities	10	17:00
10. Understanding and Managing Dysphagia	11	23:03
11. Food Texture Levels and Safe Swallowing Practices	12	25:20
12. IDDSI Level 7: Easy-to-Chew	13	26:17
13. IDDSI Level 6: Soft and Bite-Sized	14	29:06
14. IDDSI Level 5: Minced and Moist	15	31:43
15. IDDSI Level 4: Puree	16	36:04

Understanding Diarrhoea in Learning Disabilities

Diarrhoea presents unique challenges for people with learning disabilities, who may struggle to communicate their discomfort or understand the changes occurring in their bodies. Proper identification and management are essential to prevent complications and maintain dignity.

According to the Bristol Stool Chart, types 5, 6, and 7 represent loose stools, but only type 7 constitutes true diarrhoea. This visual tool helps caregivers accurately assess and communicate bowel movements to healthcare professionals, ensuring appropriate medical intervention when needed.

The causes of diarrhoea in people with learning disabilities are diverse and can include infections, dietary intolerances, medications, and underlying medical conditions. Some individuals may be at higher risk due to reduced immune function or difficulty maintaining proper hygiene practices.



The Bristol Stool Chart helps identify different types of bowel movements, with type 7 indicating true diarrhoea.

⊗ **Critical Action Required:** Any new onset of diarrhoea must be assessed by a GP, as people with learning disabilities have a higher risk of serious bowel conditions, including bowel cancer.

Causes and Initial Management of Diarrhoea

Infectious Causes

- Viral gastroenteritis
- Bacterial infections
- Parasitic infections
- Food poisoning

Medication-Related

- Antibiotics
- Laxative overuse
- Certain psychiatric medications
- New medication introductions

Dietary Factors

- Food intolerances
- Lactose intolerance
- High-fat foods
- Artificial sweeteners

Medical Conditions

- Inflammatory bowel disease
- Irritable bowel syndrome
- Coeliac disease
- Bowel cancer

When diarrhoea occurs, immediate steps should focus on preventing dehydration and seeking medical assessment. Encourage increased fluid intake with oral rehydration solutions when appropriate, monitor for signs of dehydration, and maintain detailed records of frequency, consistency, and any associated symptoms.

Remember: People with learning disabilities may not be able to express discomfort or pain effectively. Watch for behavioural changes that might indicate abdominal cramping or distress.

Dietary Management Strategies



GP Assessment First

Always consult a GP for proper diagnosis before implementing dietary changes. Some conditions require medical treatment alongside dietary modifications.



Professional Guidance

Request a dietitian referral for personalised advice, especially for complex conditions like coeliac disease or inflammatory bowel disease.



Monitor and Record

Keep detailed food and symptom diaries to identify triggers and patterns that can inform treatment decisions.

For conditions like IBS (Irritable Bowel Syndrome), dietary modifications can significantly improve symptoms. Key strategies include reducing dietary fat to less than 3g per 100g of food, avoiding spicy foods containing capsaicin (the compound that makes chillies hot), and maintaining regular meal times to provide predictability for the digestive system.

Choose naturally low-fat options such as fruits, vegetables, wholegrains, lean proteins, low-fat dairy products, beans, and pulses. Use cooking sprays instead of oils, select low-fat dressings and sauces, and encourage regular eating patterns with three main meals per day, with or without healthy snacks.

Understanding Constipation: A Critical Concern

70%

Lifetime Risk

Percentage of people with learning disabilities who will experience constipation during their lifetime

23%

Health Impact

Of those with learning disabilities who died in 2019 had constipation as a long-term condition

13%

Serious Consequences

Died from constipation-related complications such as bowel perforation or obstruction

Constipation represents a significant health concern for people with learning disabilities, with potentially life-threatening complications if left untreated. The statistics above highlight the critical importance of early recognition and appropriate management.

The higher prevalence of constipation in this population stems from multiple factors including reduced mobility, medication side effects, dietary limitations, communication difficulties, and underlying neurological conditions. Many individuals may not be able to express discomfort or understand the need for regular toileting routines.

 Constipation can lead to serious complications including faecal impaction, bowel obstruction, perforation, and even death. Early intervention is crucial.

Recognising and Responding to Constipation

Warning Signs to Monitor

Behavioural Changes

Increased agitation, aggression, or distress may indicate abdominal pain that cannot be communicated effectively.

Absence of Bowel Movements

No bowel movement for three days or more requires immediate attention and medical assessment.

Unusual Incontinence

New episodes of faecal incontinence in someone who is typically continent may indicate overflow.

Physical Signs

Blood in underwear, extended bathroom visits, or visible discomfort during toileting attempts.



Early recognition of constipation symptoms enables prompt medical intervention and prevents serious complications.

When constipation is suspected, immediate medical consultation is essential. Both constipation and diarrhoea can indicate serious underlying conditions, and people with learning disabilities face higher risks for bowel cancer and other gastrointestinal disorders. Document all symptoms, behavioural changes, and toileting patterns to provide comprehensive information to healthcare professionals.

Prevention Strategies for Constipation



Adequate Hydration

Encourage 6-8 glasses of fluid daily, including water, squash, tea, coffee, and juice. Use personalised cups or straws to make drinking more appealing. Regular fluid intake throughout the day helps maintain soft stool consistency.



Balanced Fibre Intake

Include both soluble fibre (oats, seeds, soft fruits) to soften stool and insoluble fibre (wholegrain carbohydrates, fruit skins) to promote movement. Aim for 2 portions of soluble and 1-2 portions of insoluble fibre with each meal.



Regular Movement

Encourage at least 30 minutes of daily physical activity. This can include walking, dancing, swimming, or adapted exercises suitable for the individual's abilities. Movement stimulates bowel function naturally.

Prevention remains the most effective approach to managing constipation. Creating structured routines around hydration, nutrition, and physical activity helps establish healthy bowel patterns. For individuals with swallowing difficulties or texture preferences, work with speech and language therapists and dietitians to ensure adequate fibre and fluid intake whilst maintaining safety.

Regular toileting schedules, particularly after meals when the gastrocolic reflex is strongest, can help establish routine. Ensure privacy and comfort during toileting, and allow sufficient time without rushing. Some individuals may benefit from visual schedules or social stories about healthy toileting habits.

Managing Reflux in Learning Disabilities

 Between 50-70% of people with severe and profound learning disabilities experience reflux.

Gastroesophageal reflux occurs when stomach contents flow backwards into the oesophagus, causing pain, discomfort, and potential complications such as Barrett's Oesophagus, which can lead to oesophageal cancer if left untreated. This condition is significantly more prevalent in people with learning disabilities, particularly those with severe or profound disabilities.

The higher incidence relates to factors such as reduced mobility, certain medications, anatomical differences, and difficulty maintaining optimal positioning after meals. Many individuals cannot communicate the burning sensation or discomfort associated with reflux, making recognition challenging.

Pain and Distress Burning sensation and discomfort that may manifest as behavioural changes	Eating Difficulties May lead to food refusal or reduced intake due to associated pain	Dental Problems Stomach acid causes tooth erosion and decay requiring dental intervention
Breathing Issues Aspirated stomach contents can cause respiratory problems and infections		Cancer Risk Untreated reflux can lead to Barrett's oesophagus and oesophageal cancer

Management involves both medical and lifestyle interventions. GPs can prescribe medications to reduce acid production or improve stomach emptying, adjust existing medications that may worsen reflux, and refer to specialists when necessary.

Preventing Reflux in Individuals with Learning Disabilities

Reflux can be a common and uncomfortable issue for individuals with learning disabilities, often exacerbated by communication challenges or physical limitations. Implementing practical strategies can significantly reduce symptoms and improve quality of life. This guide provides actionable advice for caregivers.

Dietary Adjustments

- Offer small, frequent meals rather than large ones.
- Identify and avoid trigger foods (e.g., fatty, spicy, acidic foods, decaffeinated and regular coffee, chocolate, alcohol, citrus based juices and fizzy drinks).
- Ensure food textures are easy to chew and swallow, reducing swallowed air.
- Encourage adequate hydration between meals, not during, to avoid overfilling the stomach.

Meal Timing & Routine

- Allow at least 2-3 hours between the last meal or snack and lying down for sleep or rest.
- Establish a consistent schedule for meals and snacks to regulate digestion.
- Avoid rushing meals; allow ample time for eating in a relaxed atmosphere.

Positioning Strategies

- Maintain an upright position during and for at least 30-60 minutes after meals.
- For sleep, consider elevating the head of the bed (if safe and appropriate, consult a professional).
- Avoid bending over or strenuous activities immediately after eating.
- Ensure clothing around the abdomen is loose and comfortable.

Environmental & Other Factors

- Create a calm, quiet, and comfortable eating environment free from distractions.
- Ensure comfortable seating that supports good posture during meals.
- Monitor any medications for side effects that might contribute to reflux.
- Regularly consult with healthcare professionals for personalized advice and management.

Understanding and Managing Dysphagia

Dysphagia, or swallowing difficulty, presents serious risks for people with learning disabilities who may struggle to understand or communicate their swallowing problems. This condition requires immediate professional assessment and ongoing monitoring to prevent aspiration pneumonia and choking incidents.

O1

Recognition of Symptoms

Watch for coughing during or after eating, wet or gurgly voice changes, food regurgitation, difficulty with specific textures, or complaints of food getting stuck. Frequent chest infections may also indicate aspiration, alongside a panicked look or colour change in their face when they swallow.

O2

Immediate GP Consultation

Any suspected swallowing difficulty requires urgent medical assessment. Do not delay seeking help, as dysphagia can quickly lead to serious complications including pneumonia and choking.

O3

Speech and Language Therapy Referral

The GP will refer to a Speech and Language Therapist (SaLT) who specialises in swallowing assessments. They will conduct comprehensive evaluations to determine safe food and fluid textures.

O4

Implementation of Recommendations


Follow SaLT recommendations precisely regarding food textures, positioning during meals, and feeding strategies. These recommendations are critical for safety and must be consistently applied.

The International Dysphagia Diet Standardisation Initiative (IDDSI) provides standardised levels for food textures and fluid thickness, ranging from Level 0 (thin fluids) to Level 7 (regular foods). Each level has specific requirements designed to reduce aspiration risk whilst maintaining nutritional adequacy.

Food Texture Levels and Safe Swallowing Practices

The IDDSI Framework

Providing a common terminology for describing food textures and drink thicknesses to improve safety for individuals with swallowing difficulties.



© The International Dysphagia Diet Standardisation Initiative 2019 @ <https://iddsi.org/framework/>
Licensed under the Creative Commons Attribution Sharealike 4.0 License <https://creativecommons.org/licenses/by-sa/4.0/legalcode>.
Derivative works extending beyond language translation are NOT PERMITTED.

Successful dysphagia management requires strict adherence to prescribed texture levels, proper positioning during meals, adequate time for eating without rushing, and regular review with the speech and language therapy team. They will prescribe a diet based off of the IDDSI framework. Never deviate from recommended textures without professional approval, as this could result in serious harm.

Support teams must receive comprehensive training on safe feeding practices, including recognition of distress signs, appropriate meal pacing, and emergency procedures. Regular reassessment ensures that texture recommendations remain appropriate as conditions change over time.


IDDSI Level 7: Easy-to-Chew

The IDDSI Level 7 Easy-to-Chew diet is specifically designed for individuals who are able to easily chew soft, bite-sized pieces of food. Critically, they can recognize when a food piece is too large and are capable of chewing softer textures and moving them around their mouth with ease.

This dietary level introduces slightly more restriction than a regular diet, emphasizing normal, everyday foods that naturally possess a softer or more tender consistency. It is well-suited for people who experience chewing and/or swallowing difficulties but can still safely manage regular foods. Examples of foods to avoid on this texture include fibrous textures like celery or pineapple, or clear seeds and pips, sticky foods like toffees or sticky mash, raw veggies, crispy foods like dry cereal or crackling, hard foods like crisps, meats which are tougher or have gristle/ fat, or small, easy-to-choke on foods like nuts or seeds.



7 EASY TO CHEW



Must be able to break food apart easily with the side of a fork or spoon

Easy to Chew foods must break apart easily and pass Fork Pressure Test!

IDDSI Fork Pressure Test

To make sure the food is soft enough, press down on the fork until the thumbnail blanches to white, then lift the fork to see that the food is completely squashed and does not regain its shape

IDDSI Level 6: Soft and Bite-Sized

Foods are soft and pre-cut into pieces no larger than 1.5cmx1.5cm. This level eliminates hard, stringy, or choking-hazard foods whilst maintaining variety.

Foods You CAN Have

- Soft cooked vegetables (mashed potato, soft carrots)
- Tender meats cut into small pieces (soft chicken, fish)
- Fruits with no skin or pips, that are soft
- Soft pasta
- Scrambled eggs
- Soft cheese
- Yogurt and puddings
- Fully softened cereal with moisture removed
- Rice moistened with sauce

Foods to AVOID

- All things from L7 avoid list, plus:
- Hard foods (nuts, raw vegetables, crusty bread)
- Stringy foods (celery, pineapple, tough meats)
- Mixed textures (soup with chunks, cereal with milk)
- Juicy foods like cucumber or melon
- Dry foods like cake without cream
- Sticky foods (peanut butter, toffee)
- Small hard items (seeds, corn kernels)
- Foods which separate into individual grains (unmoistened rice)

Preparation Tips

- Cut all foods to 1.5cm cubes or smaller
- Ensure all foods are fork tender (can be squashed with a fork and do not regain shape)
- Cook vegetables until soft
- Remove skins and seeds from fruits
- Ensure meats are tender and well-cooked

IDDSI Level 5: Minced and Moist

Foods are minced with lumps no bigger than 4mm (the size of the gap between fork prongs) and served with a thick, non-pouring sauce or gravy. The sauce should not separate from the food. This level requires minimal chewing and eliminates separate textures, providing a safe and easy-to-swallow option.

Foods You CAN Have

- Minced meat (e.g., chicken, beef) with thick gravy
- Mashed fish with sauce
- Soft, mashed vegetables (e.g., carrots, pumpkin, potato)
- Thickened soups and stews
- Yogurt and smooth puddings
- Soft, moist casseroles (minced)
- Cereal which has been moistened into a thick paste with excess liquid removed.
- Sieved/ finely mashed fruit with no skins or seeds

Foods to AVOID

- All things from L6 and L7 avoid list, plus:
- Hard, dry, or crumbly foods
- Foods with separate liquids (e.g., thin broth, watery sauces)
- Sticky foods (e.g., peanut butter, toffee)
- Foods with skins or seeds (including meats or salmon, beans and peas)
- Lumps larger than 4mm
- Stringy foods (e.g., celery, fibrous meats)

Preparation Tips

- Mince all foods to 4mm or smaller
- Add thick sauce, gravy, or custard to ensure moistness
- Ensure food holds its shape on a spoon but is easily mashed with a fork
- Avoid adding extra water as this can thin out the consistency, flavour and nutrition. Add thick gravies, custards, white sauces and other flavourful, nutritious sauces to achieve the correct consistency.

IDDSI Level 4: Puree

Foods at this level have a smooth, lump-free texture that requires no chewing. They must not be sticky and should not separate into different consistencies. The food can be spooned but cannot be poured.

Foods You CAN Have

- Smooth pureed meats (e.g., chicken, beef)
- Pureed vegetables (e.g., carrots, peas, butternut squash)
- Smooth pureed fruits (e.g., applesauce, pureed peaches)
- Thick, smooth soups and gravies
- Yogurt and smooth custards without lumps
- Pureed cereals (e.g., smooth porridge)
- Mashed potatoes

Foods to AVOID

- All foods excluded from L5, L6 and L7, plus:
- Any foods with lumps, seeds, or skins
- Hard, crunchy, or crumbly foods
- Sticky foods (e.g., thick honey, very dense peanut butter)
- Foods that separate liquid (e.g., thin soups, some pureed fruits with juice)
- Rice or any other non-smooth food
- Stringy or fibrous foods
- Foods which get a crust whilst cooking (such as a cottage pie)
- Jellies

Consistency Tests

- **Fork Drip Test:** Place food on a fork and tilt it. The food should sit in a mound or dollop and flow slowly through the gaps, leaving a coating on the fork. It should not drip continuously.
- **Spoon Tilt Test:** The food should hold its shape on a spoon when tilted sideways, but easily fall off with a slight tap or gentle shake. It should not stick to the spoon.
- **No Chewing Required:** The food should be smooth enough to swallow without any chewing.